


MEDICAL STUDENT VIRTUAL RESEARCH SYMPOSIUM GUIDELINES

Dr. Fern Tsien
Assistant Dean, Medical Student Research
Department of Genetics
LSUHSC



Who is eligible to present at the Virtual Research Symposium?

- Medical students from LSUHSC and partnering medical schools
- Summer program, MCLIN198, and Honors Program students are highly encouraged
- You are welcome to present more than one poster.
- Good practice for national and international conferences, and can be added to your resume/CV
- If co-authoring, each student presents a separate poster with each as first author).
- Student presentations will be judged and awards will be given for each category
 - 1st and 2nd year med students
 - 3rd and 4th year med students

Important deadline #1:

- Friday, September 24th
- If you plan to participate, please contact us at SoMHonorsProgram@lsuhsc.edu so we can send you notifications and guidelines

Important Deadline #2: Abstracts

- Abstracts are due on or before Monday, October 4th, 2021 at 5:00 pm!!!
- If you already turned in an abstract this summer, please resubmit it, even if there are no changes to: SoMHonorsProgram@lsuhsc.edu
- Follow the guidelines sent to you. DO NOT change the margins, font size, or font style.
- These will be sent to the judges ahead of time.

Sending the abstracts

- One-page summary of your project
- List your name and mentor's name as described in the template.
- Affiliations: department and school.
- Use only the template we provide.
- This template has the correct sized fonts and sizes we will use. Do not change the font or size!
- Follow the directions provided by Ms. Kirsten Bruno.
- Make sure your mentor approves of your abstract before you send it to us!
- When you submit your abstract in Word format, please be sure to save the file with your last name listed first. For example: **BrunoKirstenAbstract.doc**
- Send it to: SoMHonorsProgram@lsuhsc.edu

Your Name (first, middle initial, last)
Classification (High School, Undergraduate, Medical)
Name of School, City, State

Mentor's Name:
Mentor's Affiliation (LSUHSC, Tulane SOM, Xavier, Children's Hospital, etc.)

"Title of Project"

Abstract (summary of project, not to exceed one page)

Body of Abstract: Left Justified, 11 pt Arial font.

|

Katherine A. Adler
L2
LSU Health Sciences Center, New Orleans, LA

Mentor's Name: Dr. Liz Simon
LSUHSC Department of Physiology


"Circulating myomiR levels as a clinical indicator of alcohol-induced skeletal muscle dysfunction in PLWH"

There are an estimated 1.15 million people living with HIV (PLWH) in the US. The prevalence of at-risk alcohol use among PLWH is higher than in the general population. Antiretroviral therapy (ART) has significantly reduced patient mortality, and HIV infection has emerged as a chronic disease with associated comorbidities such as myopathy and insulin resistance. Impaired skeletal muscle (SKM) function and mass is a consistent predictor of mortality and contributes to a decrease in quality of life in PLWH. Chronic alcohol and HIV independently and synergistically contribute to significant SKM derangements such as atrophy, weakness, and dysfunction. Previous studies have shown that chronic alcohol exposure alters the epigenome including muscle specific microRNA (myomiR) expression, correlating with alterations in expression of myogenic genes. MicroRNAs are produced in cells and secreted actively or passively into circulation. Abundance of circulating myomiRs is a function of the regenerative and degenerative capacity of the muscle, the overall muscle mass, and tissue expression levels. Our hypothesis was that circulating myomiRs is decreased in PLWH with at-risk alcohol use, and they would correlate with a decrease in SKM mass and function.

Subjects from the LSUHSC HIV Outpatient Program were stratified into low, mid, and high drinkers based on timeline follow back (TLFB) and corresponding AUDIT scores. Circulating myomiR levels were determined and correlated to measures of AUD severity, hand grip strength, 4-meter walk test, and lean mass.

The muscle-specific miRNAs 206 and 133a expression were significantly increased in individuals with mid- and high-drinking. Copy number calculations of these myomiRs revealed they were positively correlated with TLFB. Sex differentially modulated the relationship, with miR-206 positively correlating with hand grip strength in males.

Contrary to our hypothesis, circulating myomiRs were increased in individuals with at-risk alcohol use. This may be due to alcohol-mediated damage or inflammation in SKM tissue. Confounding variables including high BMI, high fat mass, and low physical activity in low-drinking cohort may have impacted circulating myomiRs and further studies will investigate correcting for these variables and using a composite myomiR score to correlate with SKM function.




What is
wrong
with this
abstract?

Mechanisms Underlying the Sleep Promoting Effect of Cherry Juice Standardized to its Proanthocyanidin Content

Previous studies have shown that tryptophan, melatonin, and proanthocyanidin within cherry juice may play essential roles in promoting sleep. This study utilizes cherry juice standardized to its proanthocyanidin content and tests its effectiveness as a treatment for insomnia, a common health problem in the elderly. Ten participants with insomnia complete two treatment periods (cherry juice and placebo juice), 2 weeks each, separated by a 2 week washout period. Each day the participants consume 8 ounces of juice in the morning and again 1-2 hours before bedtime. Overnight polysomnography (PSG) is used at the end of each treatment period to evaluate sleep architecture such as the distribution of sleep stages, sleep latency and state transitions. Blood samples are also taken to measure serum concentrations of free tryptophan and kynurenine in order to investigate a possible mechanism of action. Questionnaires are given before and after each two week treatment period for comparison of each treatment's effects. This study is still ongoing and data analysis will be performed upon its completion.



Important Deadline #3: Posters

- Posters are due Wednesday,
October 13th at 5:00 pm!
- 

Preparing the posters 1

- **First and most important:** make sure that your mentor approves of the information that will be presented in the poster.
- **Second most important:** Your name should go first, your mentor's name last, and everyone else who helped you (other students, post-docs, etc.) in the middle. Make sure not to leave out anyone who helped you!

Preparing the posters 2

- Make sure that you understand everything you write on the poster. You should be able to explain your project to the judges.
- In general, try to keep text towards the outside and figures and tables in the center.
- The abstract is not necessary for the poster.

Preparing the posters 3

- Use the Power Point poster template sent to you by Ms. Kirsten Bruno (not your friend's or past interns) with the proper logos.
- These correspond to your mentor's affiliation and the Summer Program funding source.
- The logos on your poster may differ from the ones on your lab mates! Do not change them!
- Use at least a 24 point font size so the text will be visible from 3 feet away.
- Feel free to adjust the box sizes and headings depending on the amount of text or figures you have.
- The poster template are already set to 34 x 44 in.

Preparing the posters 4

- Use any color you want to. Express yourself!
Exceptions:
 - Black or deep blue for background of entire poster.
 - Image enlarged to cover the entire background.
- Spell out any acronyms the first time you use them. People outside of your lab may not know what “SIV” or “FSHD” is.
- Refer to guidelines sent to you.

Once your poster is done:

- Save it as a PPT *and* PDF file.
- When you submit your poster, be sure to save the file with your last name listed first. For example: **BrunoKirstenPoster.pptx**
- Send it to: SoMHonorsProgram@lsuhsc.edu
- Let us know in advance if you want us to print one for your lab and an extra one for your school.
- **Posters are due Wednesday, October 13th at 5:00 !!!**

What is wrong with this poster?

Effect of Gain-of-Function Mutant Rb on the Sphere-Forming Ability of Cell Lines

Your name goes first, Graduate students and post-docs that helped you, Mentor is last
Mentor's department and University



PATRICK F. TAYLOR
FOUNDATION



Health
Sciences
Center

Abstract

Osteosarcoma, the most common bone cancer, is the second highest cause of cancer-related death in children and adolescents. Approximately 90% of cases show micro-metastasis at diagnosis, making systematic chemotherapy the first choice of treatment. Despite intensive chemotherapy, the survival rate for high-grade osteosarcomas remains at only 50-80%. This persistence is mainly due to the ability of osteosarcoma cells to metastasize and develop resistance to therapy. Increasing evidence suggests that cancer stem cells (CSCs) or tumor initiating cells (TICs) are responsible for these properties and that the inadequacy of current treatments may be result from the inability to target CSCs or TICs in osteosarcoma. Recently, we have demonstrated that small populations of osteosarcoma cells can grow and form spheres in both serum- and anchorage-independent manners. Importantly, as few as 200 cells from these spheres efficiently initiated osteosarcoma in tumor transplantation models. These results suggest that spheres are enriched with osteosarcoma CSCs. Approximately 50% of osteosarcomas contain alterations in the tumor suppressor p53 gene. Many hotspot p53 mutants show oncogenic functions by their gain-of-function phenotypes such as increased transformation, metastasis and drug resistance, which can not be explained simply due to loss of wild-type p53 function. Our preliminary results indicate that downregulation of a gain-of-function mutant p53R175H/72R results in a dramatic reduction of sphere-forming ability of an osteosarcoma cell line expressing p53R172H. These results suggest that mutant p53 gain-of-function is involved in sphere-forming ability and possibly CSC-like properties of osteosarcoma. However, the exact molecular mechanisms which contribute to sphere formation and CSC-like properties and the involvement of mutant p53 in these cellular phenotypes remain unclear.

Our long-term goal is to identify the molecular mechanism underlying the CSC-like properties of osteosarcoma. *The objective of this study* is to investigate the effects of several hotspot p53 mutants on the sphere-forming ability of human osteosarcoma cell lines. *Our hypothesis* is that gain-of-function p53 mutants increase the sphere-forming ability of osteosarcoma cells. To test our hypothesis, we first characterized the sphere-forming ability of several available human osteosarcoma cell lines, such as U2OS (p53 wild-type), SJS1A1 (p53 wild-type), MG63 (p53-null), Saos-2 (p53-null), and KHOS (p53R156P). We found that U2OS and MG63 cell lines did not show any sphere formation when 500 cells were tested for 2 weeks of culturing in sphere-specific conditions. These results may suggest that the presence of wild-type p53 is not crucial for the sphere formation. Assays for other cell lines are on-going. We next infected MG63 cells with retroviral vectors encoding p53R175H/72P, p53R175H/72R, p53R248W/72P, p53R248W/72R, p53R273H/72P, and p53R273H/72R to establish MG63 subcell lines expressing several gain-of-function p53 mutants together with different p53 codon 72 single nucleotide polymorphisms (SNPs), since the SNP is shown to affect colony-forming ability of human cancer cell lines. Sphere formation assays using these subcell lines are underway and all results will be presented. Completion of our study will provide a better understanding of the role of gain-of-function mutant p53 in sphere-forming ability of osteosarcoma as well as useful information to dissect the molecular mechanism of CSC-like properties of osteosarcoma.

Introduction

Osteosarcoma is a devastating disease in children and young adults. In approximately 90% of osteosarcoma cases, micro-metastases are present during diagnosis, making chemotherapy the first choice of treatment. Despite intensive chemotherapy, the survival rate for high-grade osteosarcomas remains at only 50-80%. This persistence is mainly due to the ability of osteosarcoma cells to metastasize and develop resistance to therapy. Increasing evidence suggests that cancer stem cells (CSCs) or tumor initiating cells (TICs) are responsible for the metastatic and drug-resistant properties of cancer cells and that the inadequacy of current treatments for high grade osteosarcoma may result from the inability to target osteosarcoma CSCs. CSCs represent a small fraction of a tumor's cellular population and have the ability to generate new tumors identical in cellular composition to the tumor of origin. CSCs possess the abilities of anchorage-independent, serum-independent cell growth (sphere formation), tumor initiation, self-renewability, and multilineage differentiation, as well as properties of high metastatic potential and drug resistance. We have recently reported that small number of osteosarcoma cells form spheres and these spheres are enriched with cells having CSC-like properties such as high metastatic and drug resistant properties. However, the molecular mechanism that regulates CSC-like properties of osteosarcoma remains unclear.

Cancer can arise through alterations to genes that regulate cell proliferation, apoptosis, and senescence. The tumor suppressor p53, one of the key guardians of these events, exerts its functions through transactivating numerous downstream targets. Tumor suppressor p53 has a single nucleotide polymorphism (SNP) at codon 72 which is either proline (P) or arginine (R). Recent studies have shown that the 72R form is more potent in its ability to induce apoptosis compared to the 72P form. In addition to the polymorphism, mutations in the p53 gene affect the p53 activity. Mutations in the DNA binding domain attenuate the function of p53 as a transcription factor, thereby losing its tumor suppressor activity. The importance of p53 mutation is emphasized by the clinical observation that the p53 gene is mutated in more than 50% of tumors. Mutations in the p53 gene are also observed in approximately 70% of patients with Li-Fraumeni syndrome (LFS), a human familial cancer-prone disease. LFS is characterized by early onset of various types of tumors, including osteosarcoma. Several missense mutations such as R175H, R248W, and R273H are the hotspot mutations in sporadic cancer as well as the germline of LFS patients. These p53 mutants show oncogenic functions by their gain-of-function phenotypes such as increased transformation, metastasis, and drug resistance, which can not be explained simply by loss of wild-type p53 function. The molecular mechanisms underlying the gain-of-function activities and if the codon 72 SNP affects the mutant p53's gain-of-function activities remain unclear. Further, although the gain-of-function phenotypes are similar to those of CSCs, the contributions of mutant p53 to the CSC-like properties are also unknown.

Methods and Materials

Cell lines. Human osteosarcoma cell lines U2OS, SJS1A1, Saos-2, MG-63, and KHOS/HP were purchased from American Type Culture Collection (ATCC, Manassas, VA).

Sphere culture. Cells were counted by trypan-blue staining (Sigma Biochemicals), and live cells (five per well) were plated on a 96-well ultra-low attachment plate (Corning Inc., Corning, NY, USA) in sphere-specific media consisting of DMEM/F12, progesterone (10 nM), putrescine (50 µM), insulin (12.5 µg/ml), transferrin (12.5 µg/ml), sodium selenite (12.5 ng/ml, Sigma Biochemicals), murine EGF (10 ng/ml), and murine bFGF (40 ng/ml, Pepro Tech, Rocky Hill, NJ, USA). Cells were maintained for 10–14 days and fresh aliquots of EGF and bFGF were added three times a week. Sphere formation was observed daily using under a phase-contrast microscopy (Nikon Eclipse TS100).

Western blotting. MG-63 cells infected with retroviral vectors encoding control empty or a mutant p53 (R175H, R248W, or R273H) with a codon 72 polymorphism were lysed with RIPA buffer (50 mM Tris-HCl pH 7.6, 150 mM NaCl, 1 mM EDTA, 1% sodium deoxycholate, 0.1% Triton X-100, 0.1% SDS) supplemented with protease inhibitor cocktail (Roche) (1 mM phenylmethylsulfonyl fluoride (PMSF), 0.2 mM sodium orthovanadate, and 100 mM sodium fluoride). Whole cell extracts were separated by SDS-PAGE and transferred onto PVDF membranes (GE Healthcare BioSciences). After blocking with 5% non-fat milk in 1 x Tris-buffered saline (TBS) with 0.1% Tween-20 (TBS-T), blots were incubated with anti-human p53 (DO1, Santa Cruz) and control vinculin (Fitzgerald), followed by the incubation with secondary antibodies conjugated with horseradish peroxidase (Santa Cruz). To visualize signals, Super Signal West Dura Chemiluminescent substrates (Pierce Biotechnology) were used according to manufacture instructions. The signals were detected using a Biorad *Verse Doc* detection system (Biorad).

Sphere culture. Cells were counted by trypan-blue staining (Sigma Biochemicals), and live cells (five per well) were plated on a 96-well ultra-low attachment plate (Corning Inc., Corning, NY, USA) in sphere-specific media consisting of DMEM/F12, progesterone (10 nM), putrescine (50 µM), insulin (12.5 µg/ml), transferrin (12.5 µg/ml), sodium selenite (12.5 ng/ml, Sigma Biochemicals), murine EGF (10 ng/ml), and murine bFGF (10 ng/ml, Pepro Tech, Rocky Hill, NJ, USA). Cells were maintained for 10–14 days and fresh aliquots of EGF and bFGF were added three times a week. Sphere formation was observed daily using under a phase-contrast microscopy (Nikon Eclipse TS100).

Western blotting. MG-63 cells infected with retroviral vectors encoding control empty or a mutant p53 (R175H, R248W, or R273H) with a codon 72 polymorphism were lysed with RIPA buffer (50 mM Tris-HCl pH 7.6, 150 mM NaCl, 1 mM EDTA, 1% sodium deoxycholate, 0.1% Triton X-100, 0.1% SDS) supplemented with protease inhibitor cocktail (Roche) (1 mM phenylmethylsulfonyl fluoride (PMSF), 0.2 mM sodium orthovanadate, and 100 mM sodium fluoride). Whole cell extracts were separated by SDS-PAGE and transferred onto PVDF membranes (GE Healthcare BioSciences). After blocking with 5% non-fat milk in 1 x Tris-buffered saline (TBS) with 0.1% Tween-20 (TBS-T), blots were incubated with anti-human p53 (DO1, Santa Cruz) and control vinculin (Fitzgerald), followed by the incubation with secondary antibodies conjugated with horseradish peroxidase (Santa Cruz). To visualize signals, Super Signal West Dura Chemiluminescent substrates (Pierce Biotechnology) were used according to manufacture instructions. The signals were detected using a Biorad *Verse Doc* detection system (Biorad).

Figure 1

Western blotting for mutant p53




Figure 2. Western blotting. MG-63 cells were infected with mutant p53-encoding retroviral vectors to generate subcell lines expressing mutant p53. Cells were lysed in RIPA buffer and whole cell extracts were subjected to western blotting for p53 (DO1 antibody) and loading control vinculin. U2OS having wild-type p53 was used as a positive control.

Results

Our long-term goal is to identify the molecular mechanism underlying the CSC-like properties of osteosarcoma. *The objective of this study* is to investigate the effects of several hotspot p53 mutants on the sphere-forming ability of human osteosarcoma cell lines. *Our hypothesis* is that gain-of-function p53 mutants increase the sphere-forming ability of osteosarcoma cells. To test our hypothesis, we first characterized the sphere-forming ability of several available human osteosarcoma cell lines, such as U2OS (p53 wild-type), SJS1A1 (p53 wild-type), MG63 (p53-null), Saos-2 (p53-null), and KHOS (p53R156P). We found that U2OS and MG63 cell lines did not show any sphere formation when 500 cells were tested for 2 weeks of culturing in sphere-specific conditions. These results may suggest that the presence of wild-type p53 is not crucial for the sphere formation. Assays for other cell lines are on-going. We next infected MG63 cells with retroviral vectors encoding p53R175H/72P, p53R175H/72R, p53R248W/72P, p53R248W/72R, p53R273H/72P, and p53R273H/72R to establish MG63 subcell lines expressing several gain-of-function p53 mutants together with different p53 codon 72 single nucleotide polymorphisms (SNPs), since the SNP is shown to affect colony-forming ability of human cancer cell lines. Sphere formation assays using these subcell lines are underway and all results will be presented. Completion of our study will provide a better understanding of the role of gain-of-function mutant p53 in sphere-forming ability of osteosarcoma as well as useful information to dissect the molecular mechanism of CSC-like properties of osteosarcoma.

Table 1

Table 1. Results of sphere formation assays

| Cell lines | p53 status | Cell# examined | # of spheres formed | % sphere formation |
|----------------|------------|----------------|---------------------|--------------------|
| U2OS | wild-type | 480 | 0 | 0 |
| SJS1A1 | wild-type | 480 | 1 | 0 |
| Saos-2 | null | 480 | 318 | 66.3 |
| MG63 | null | 480 | 0 | 0 |
| MG63 R175H/72P | R175H/72P | 480 | 84 | 17.5 |
| MG63 R175H/72R | R175H/72R | 480 | 160 | 33.3 |
| MG63 R248W/72P | R248W/72P | 480 | 217 | 45.2 |
| MG63 R248W/72R | R248W/72R | 480 | 144 | 30.0 |
| MG63 R273H/72P | R273H/72P | 480 | 112 | 23.3 |
| MG63 R273H/72R | R273H/72R | 480 | 136 | 28.3 |
| KHOS | R156P | 480 | 112 | 23.3 |

Conclusions

Conclusions

- Spheres vary in size and rate of growth in different osteosarcoma cell lines.
- The presence or absence of wild-type p53 does not have any effects on the sphere-forming ability of osteosarcoma cell lines.
- The presence of mutant p53 does enhance the sphere formation of osteosarcoma cells.
- The effects of p53 codon 72 polymorphisms vary in different p53 mutants.
- All p53 mutants confer osteosarcoma cells with sphere-forming abilities.

Future directions

- Examine the effects of mutant p53 on other CSC-like properties such as tumor initiating ability, self-renewal, metastatic potential, and drug resistance.
- Examine the effects of mutant p53 down-modulation in various osteosarcoma cell lines carrying mutant p53.
- Identify genes that regulate sphere-forming ability and CSC-like properties of osteosarcoma cells.

Example of a better poster



AXIN2 Gene Instability In Colon Cancer

Summer Student (you), People who helped you, mentor
Mentor's department and University



Abstract

Colon cancer is one of the most prevalent and fatal cancers in the world. In the United States, 10% of all cancer patients have colon cancer. The disease begins when adenomatous polyps, fleshy growths that line up on the inside of the colon, become cancerous. Colonoscopy is often performed to detect these polyps. Regular testing after the age of 40 can drastically reduce the risk of developing colon cancer.

The AXIN2 gene, located in the region of 17q23-q25, is a gene of interest due to its interaction with the Adenomatous polyposis coli (APC) gene in the Wnt signaling pathway and its association with colon cancer with defective mismatch repair. Mutations in the Adenomatous polyposis coli (APC) gene have been found in about 85% of colon cancer patients. However, not much is currently known about the role of AXIN2 in colon cancer development. By conducting research on AXIN2, researchers are hoping that this gene may assist in distinguishing different subgroups of colon cancer. For this project, we analyzed two colon cancer cell lines to determine their karyotypic differences and for any 17q23-q25 region abnormalities.

The majority of the metaphase cells from both of the colon cancer cell lines analyzed were aneuploid, with one cell line (SW480) having a dramatically higher number of chromosomes reaching hypertetraploidy (103 chromosomes). In addition, the SW480 cell line contained some metaphase cells with an extra copy of chromosome 17 with amplification of the 17q23-25 region. This is the gene location of AXIN2, indicating the possibility of AXIN2 over-expression leading to the colon cancer in this cell line.

Introduction

The colon is the last portion of the large intestine, which also includes the rectum. Colorectal cancer (CRC), also known as colon cancer, is the third most common cancer in the world and the second most fatal cancer in the Western hemisphere. It is reported that approximately 655,000 people worldwide die from this disease every year. It usually arises from adenomatous polyps that line the inside of the colon. Mutations in certain genes have been associated with this disease.

One significant gene known to cause CRC is the adenomatous polyposis coli gene (APC). The APC gene is located on the chromosome 5 between positions 21 and 22. Its normal function is to provide instructions for the creation of the APC protein, which helps control how and when a cell should divide. Mutations in this tumor suppressor gene can cause CRC, gastric (stomach) cancer, and Turcot syndrome. Approximately 85% of the people who have colon cancer have a mutation in the APC gene. If a person inherits just one defective copy of the gene from one of their parents, then he or she is almost guaranteed that they will develop colon cancer by the age of 40.

A gene that the APC interacts with is the relatively unknown AXIN2 gene, the focus of this project. Located on chromosome 17 between positions 23 and 24, this gene's protein, Axin2, is presumably very important in the regulation of beta-catenin, which is also a function of the APC gene. Since the APC gene and AXIN2 gene interact in the same pathway, it is believed that a mutation to either gene can affect the other gene. About 30% of the people with colon cancer with defective mismatch repair (the mechanism to correct DNA replication errors) have a mutated AXIN2 gene. The region containing the gene shows loss of heterozygosity in breast cancer, neuroblastoma, and other cancers and tumors. Deletions or mutations in this gene can result in truncated proteins which are most likely inactive. There is a possibility that somatic inactivating mutations in AXIN2 can deregulate beta catenin, and therefore, AXIN2 may be tumor suppressor gene.

Colon Cancer Symptoms

- Constipation
- Vomiting
- Stomach cramps
- Thin stool
- Diarrhea
- Unexplained Weight loss
- Hematochezia (Blood in stool)

Figure 1

The AXIN2 gene is located on Chromosome 17 on the q arm (long arm) between positions 23 and 24. The gene spans about 35 kbp and 843 amino acids.




Figure 2

The Four Stages of Colon Cancer




Figure 3





Figure 4

G-banded Metaphases From Colon Cancer Cell lines



Methods and Materials

Samples and Culture Conditions:

Two colon cancer lines were obtained from human patients. The Sw48 cell line was obtained from an 82 year old female and the SW480 cell line was obtained from a 50 year old male. The cells were grown in DMEM with 10% Fetal Bovine Serum (FBS) and 1% penicillin under normal culturing conditions.

Chromosome Preparation:

For solid staining and G-banding, cells were harvested in exponential phase, incubated with colcemid, treated with a KCL hypotonic, and fixed two times with methanol and acetic acid. For solid staining, the cells were dropped onto slides and stained with Giemsa. For G-banding, the cells were dropped onto slides, followed by a short incubation in a trypsin solution prior to staining with Giemsa.


Results

Ploidy of Human Colon Cancer Cell Lines


| | Sw48 | Sw480 |
|------------------------------|----------|---------|
| Total # of cells analyzed | 35 | 20 |
| Diploidy = 46 (Normal #) (%) | 2 (6%) | 0 (0%) |
| Hyperdiploidy 47-57 (%) | 33 (94%) | 6 (30%) |
| Hypotriploidy 58-68 (%) | 0 (0%) | 8 (40%) |
| Triploidy = 69 (%) | 0 (0%) | 0 (0%) |
| Hypertriploidy 70-80 (%) | 0 (0%) | 1 (5%) |
| Hypotetraploidy 81-91 (%) | 0 (0%) | 4 (20%) |
| Tetraploidy 92 (%) | 0 (0%) | 0 (0%) |
| Hypertriploidy 93-103 (%) | 0 (0%) | 1 (5%) |

The table to the right shows the frequency of different ploidy levels in the Sw48 and Sw480 colon cancer cell line.

Sw48 Cell



Sw480 Cell



G-Banded Karyotypes Representative of Colon Cancer Cell Lines. The Red Arrows indicate abnormalities.

49, XXX, Dcl (1), (p31), -3, +7, +9, inv (14) (q11q22), +18, +21

57, X,Y, +der X, iso (1q), +2, iso (3q), -4,-6,-8, +10,+11,+12,+13,+15,+17,+add (17) (q23), +21,+22

Conclusions and Future Directions

When compared to normal human diploid cells, the majority of the cells from the Sw48 cell line were hyperdiploids ranging from a total of 47 to 57 chromosomes per cell, while the Sw480 cell line had a wide range of total chromosome number ranging from hyperdiploidy to hypertetraploidy (up to 103 chromosomes). Our results had many similarities with published literature on these cell lines. For example, both previously published and our analysis of sw40 showed the presence of some diploid cells as well as some hyperdiploidy, with an extra chromosome 7 in common.

The sw480 cell line was much more unstable in both studies, with common abnormalities including a missing Y, an extra X abnormal X chromosome, isochromosome 3q, and trisomy 13, 21, and 22. The previous report found one extra chromosome 17. However, our results show four 17 chromosomes, with one of them containing additional genetic material at the q23-ter, the critical region of the AXIN2 gene. Fluorescence *in situ* hybridization (FISH), RNA, and protein analyses should be performed to determine the extent of AXIN2 amplification in the Sw480 cell line.

Due to the nature of these immortalized cell lines, chromosome abnormalities are acquired with increased cell proliferation. *In vitro* studies such as this one can help to give an idea of what can occur *in vivo*. More cancer cell lines should be analyzed in order to find genetic differences between the various types of colon cancer.

Geriatric Depression Scale Scores Correlate With Changes in the Oral Microbiota and Abundances of Opportunistic Pathogens in HIV Positive Individuals



William Byerley, Eugene Blanchard, Vincent Maffei
 Meng Luo PhD, David Walsh MD, Christopher Taylor PhD
 Department of Microbiology, Immunology, and Parasitology
 Louisiana State University Health Sciences Center New Orleans



Introduction

Several studies provide evidence that there is a link between depression/mental illness and microbial community, particularly the gut. However, there has been little research into the link between depression and the population of oral microbiota, especially for individuals with Human Immunodeficiency Virus (HIV).

We hypothesized that there will be a significant difference in the oral microbiota of individuals with depression and those without depression. Furthermore, we hypothesized that HIV positive patients with depression will show a higher abundance of opportunistic pathogens than patients without depression. Similarly, the abundance is expected to be higher for HIV negative patients with depression when compared to those without depression.

Sample Demographic

| | 0-5 | 6-15 |
|-----------------|------------|------------|
| n | 51 | 25 |
| Male | 24 (47.1%) | 10 (40%) |
| Female | 13 (25.5%) | 15 (60%) |
| HIV (+) | 37 (72.5%) | 20 (80%) |
| HIV (-) | 14 (27.5%) | 5 (20%) |
| Age (yr) | 50.4 ± 5.5 | 58.4 ± 7.3 |
| White | 19 (37.3%) | 6 (24%) |
| Black | 23 (45.1%) | 19 (76%) |
| American Indian | 1 (2%) | 0 (0%) |
| Smoking | 26 (51%) | 17 (68%) |
| Alcohol | 51 (100%) | 23 (92%) |
| Drug Abuse | 32 (62.7%) | 20 (80%) |

Table 1. Sample survey ranged into the categories of those with a GDS score of 0-5, and a GDS score of 6-15. The categories for Sex, Male, Female, etc. are reported for the number of samples that meet the category criteria listed by the percentage.

Alpha Diversity




Figure 1. Overlaid box plot, "Shannon's H'", of the groups: HIV+ (GDS 0-5) and HIV+ (GDS 6-15), HIV- (GDS 0-5) and HIV- (GDS 6-15). The HIV+ (GDS 0-5) group has a significantly higher median diversity than the other three groups. However, we did not expect the other comparisons to come back as not statistically significant.

Unifrac Pcoa Plot




Figure 2. Weighted Unifrac plot.

Taxonomic Summary




Figure 3. Bar chart showing relative abundance of 22 bacterial taxa in HIV+ (GDS 0-5), HIV+ (GDS 6-15), HIV- (GDS 0-5), and HIV- (GDS 6-15) groups. The taxa are ordered by their relative abundance in the HIV+ (GDS 0-5) group. The taxa are color-coded by their relative abundance in the HIV+ (GDS 0-5) group. The taxa are ordered by their relative abundance in the HIV+ (GDS 0-5) group.

| Taxonomy | Total | HIV+ (GDS 0-5) | HIV+ (GDS 6-15) | HIV- (GDS 0-5) | HIV- (GDS 6-15) |
|-----------------------|-------|----------------|-----------------|----------------|-----------------|
| Prevotella intermedia | 0.8% | 0.8% | 0.8% | 1.7% | 0.8% |
| Prevotella nigrescens | 0.8% | 0.8% | 1.0% | 0.7% | 0.8% |
| Prevotella pallens | 1.0% | 1.0% | 1.0% | 1.0% | 1.0% |
| Streptococcus sp. | 1.0% | 0.8% | 0.8% | 1.7% | 1.0% |

Table 2. Out of the 22 opportunistic pathogens studied only 5 were more abundant in the HIV+ (GDS 0-5) group than in the other three groups. The taxa in blue indicate a relationship with GDS scores. The taxa in yellow indicate a relationship with HIV status. The taxa in red indicate a relationship with both HIV status and GDS scores. The taxa in green indicate a relationship with HIV status only. The taxa in orange indicate a relationship with GDS scores only. The taxa in purple indicate a relationship with both HIV status and GDS scores.

Conclusions

- Diversity (Shannon's H') shows an association with GDS scores.
- The majority of subjects with higher GDS scores were HIV positive.
- Of the 22 pathogens investigated, only Streptococcus sp., Prevotella intermedia, and Prevotella nigrescens demonstrated a relationship with GDS scores and not HIV status.
- Alpha Diversity only showed statistical significance for the groups HIV+ (GDS 0-5), and HIV- (GDS 0-5).

“Unexpected Results from Hereditary Cancer Panel Genetic Testing: Do Duplications of MMR Genes Matter?”

Sophia Turner¹, Alix D’Angelo, MGC, CGC^{1,2}.

¹Louisiana State University Health Sciences Center, Department of Genetics

²University Medical Center New Orleans, Cancer Center.




Introduction

- Lynch syndrome (LS), is the most common form of hereditary colorectal cancer (up to 82% lifetime risk), and also increases the lifetime risk of a variety of cancers, including endometrial (up to 60%), ovarian (up to 24%), gastric, small bowel, hepatobiliary tract, pancreatic, urinary tract, brain and skin neoplasms.^{1,2}
- LS is inherited in an autosomal dominant pattern and caused by heterozygous germline mutations in one of five genes: *MLH1*, *MSH2*, *MSH6*, *PMS2*, and *EPCAM*.⁴
- MLH1*, *MSH2*, *MSH6* and *PMS2* are known as mismatch repair (MMR) genes, which play a major role in DNA repair due to replication errors.³ *EPCAM* is not an MMR gene, however, it impacts the expression of *MSH2*.⁴
- Identifying individuals with LS is crucial, because increased surveillance and preventative surgical options are available.⁷
- We present two patients who were referred to the Genetic Counseling clinic at University Medical Center. Interestingly, both patients met *BRCA1/2* genetic testing criteria but were found to have a duplication of an MMR gene.
- A literature search was performed to determine whether these duplications may be of clinical significance, and therefore impact patient management.


Hereditary Cancer Panels



Genetic Test Results




Patient #1




Patient #1 is a 57 year-old post-menopausal female referred to Genetics clinic due to her personal and family history of cancer. She was diagnosed with stage IB ERPR+ HER2- invasive ductal carcinoma of the left breast at 50 years-old. She underwent left mastectomy and 6 weeks of adjuvant radiation therapy. She has a history of smoking tobacco (1/2 pack per day since she was 20 years-old). Details of the family history are available in the pedigree above.

Patient #2



Patient #2 is a 31 year-old pre-menopausal, nulliparous, unaffected female referred to Genetics due to her family history of cancer. Details of the family history are available in the pedigree above.

Variant Classification Scheme



Discussion

- Next-generation sequencing technology has drastically transformed the genetic testing paradigm, particularly in the hereditary cancer specialty.^{8,9} However, this testing has also led to an increase in inconclusive and unexpected results.¹⁰
- Despite the patients' histories of breast +/- ovarian cancer, both were found to have a whole gene duplication of an MMR gene (*MSH2*, *PMS2*). The families presented in this report do not meet Amsterdam II criteria, however, they are suspicious of hereditary forms of cancer considering the types of cancers, ages at diagnosis and number of affected relatives in the families.
- A recent study of 528 individuals who have a mutation in one of the MMR genes shows an interesting correlation with breast cancer. Among these individuals, 23.5% had breast cancer (compared to 35.2% who had colorectal cancer and 25.8% who had endometrial cancer), noting that breast cancer was nearly as prevalent. Additionally, individuals who had mutations in *PMS2* or *MSH6* were more likely to meet NCCN guidelines for *BRCA1/2* testing (not Lynch syndrome) than *MLH1* and *MSH2* carriers.¹¹ This study suggests that the presentation of our patients may be part of the LS phenotypic spectrum.
- Many different types of mutations in the MMR genes are known to be pathogenic, including missense, nonsense, deletions and partial duplications.¹²⁻¹⁷ For example, in a report of two individuals with personal and family histories of early- and late-onset colorectal, endometrial and other cancers, across 7-14 of *MSH2* were duplicated. While the families did not meet Amsterdam II criteria, several tumors were confirmed to have high microsatellite instability, which combined with the presentation of these patients confirmed that the duplication was responsible.¹²
- Unfortunately, evidence of whether whole MMR gene duplications are pathogenic is limited, and they are currently classified as variants of unknown significance (VUS). Pathogenic whole gene duplications have been observed in another gene that is associated with hereditary colorectal cancer/polypsis, *GREM1*.¹⁴⁻¹⁶ However, *GREM1* and MMR protein products serve very different functions. Additionally, there are no families that meet Amsterdam II criteria with whole MMR gene duplications that have been reported in literature, to our knowledge.
- Follow-up testing, including chromosomal microarray may be beneficial for our patients in order to further evaluate the size and location of the duplications. Further family and molecular studies are necessary to reclassify these variants, as this may have a dramatic impact on the management of patients and their families.

Influenza Vaccination Program Requirements of Healthcare Personnel in Louisiana Hospitals

names
LSUHSC-NOLA, Department of Pediatrics, Division of Infectious Diseases and Children's Hospital, New Orleans

Introduction

- Influenza virus causes 24,000 annual deaths in the U.S. Every year 450,000 to 900,000 Louisiana residents are infected and 800 die.
- To prevent high morbidity and mortality, annual vaccination of patients and healthcare personnel (HCP) is recommended. Yet, the vaccination coverage of U.S. HCP in 2010 was only 60%.
- In response, the Centers for Disease Control and Prevention (CDC) is demanding that vaccination rates improve to 90% by 2020, and various Medical Societies are recommending mandatory vaccination programs (i.e., requirement for employment).
- To improve influenza vaccination coverage of HCP in Louisiana hospitals we must first understand what is being done, what is effective and what is ineffective.

Objectives


1. To determine influenza vaccination requirements and policies among hospitals in Louisiana, including the prevalence of mandatory requirements and consequences for declination
2. To correlate specific requirements with vaccination rates achieved, and to identify interventions that may increase vaccination rates

Methods

- A survey was sent to all 256 hospitals in Louisiana (under 193 organizations) identified in the Directory of the Louisiana Hospital Association.
- The survey contained questions on type of hospital, patient population served, components of the vaccination program and their estimated vaccination rate.
- Data was inputted into an Excel sheet and analyzed for components that influenced vaccination rates.
- Univariate analysis of categorical data compared the median vaccination rate between hospitals with or without a specific variable using the non-parametric Mann-Whitney test.
- The effect of continuous variables on the vaccination rate was analyzed with regression analysis using the non-parametric Spearman r.
- A p Value of <0.05 was considered significant.

Results: Hospitals Responding

- In the first 4 weeks, 49 (25%) of the 193 administrations responded with a statewide distribution (Figure 1).




Results: Main Responses

- Most hospitals were private for profit (51%), private non-profit (35%), and public (14%); 22% were teaching and 51% were accredited by The Joint Commission.
- The median number of beds was 60 with a range of 10 – 800.
- All hospitals had a flu vaccination program; 33% had voluntary vaccination and 67% required a formal declination. No hospital demanded vaccination as a requirement of employment.
- All hospitals offered free vaccines; 27% met all CDC recommended activities for vaccination but 73% did not meet all CDC recommendations.
- 24% of hospitals enforced consequences to HCP declining vaccination while 76% had no consequences; the most common consequence was a requirement to wear a mask on patient contact.
- The median vaccination rate reported by the responding hospitals was 61%, with a range from 12 - 98%.

Results: Correlates of Vaccination

| Factors Positively Associated with Vaccination Rates | | | | | | |
|--|---------------|----|-------------------|-------------|-------|---------|
| Survey Questions | No. Responses | % | Not Present | Present | Ratio | p Value |
| | | | Median (25%, 75%) | | | |
| Hospital Type | | | | | | |
| Private | 18 | 37 | 55 (45, 72) | 73 (58, 84) | 1.33 | 0.02 |
| Acute Care | 28 | 58 | 50 (45, 72) | 70 (57, 81) | 1.40 | 0.02 |
| High-Risk Patient Type | | | | | | |
| Children | 29 | 59 | 50 (45, 71) | 70 (56, 85) | 1.40 | 0.02 |
| Pregnant Women | 23 | 47 | 51 (45, 71) | 72 (60, 85) | 1.41 | 0.004 |
| Intensive Care | 26 | 53 | 50 (42, 70) | 71 (57, 85) | 1.42 | 0.006 |
| Number of Beds | | | | | | |
| 0 - 99 | 26 | 53 | | 50 (45, 71) | 0.694 | 0.0006 |
| 100 - 299 | 12 | 24 | | 70 (56, 80) | 1.186 | |
| ≥ 300 | 8 | 16 | | 85 (61, 92) | 1.466 | |
| Vaccination Program | | | | | | |
| Voluntary | 16 | 33 | 71 (52, 85) | 52 (40, 57) | 0.73 | 0.001 |
| Declination Required | 33 | 67 | 52 (40, 57) | 71 (52, 85) | 1.37 | 0.001 |
| Vaccine Administration | | | | | | |
| Common areas | 31 | 63 | 48 (37, 52) | 70 (59, 83) | 1.46 | 0.001 |
| Nights/Weekends | 38 | 78 | 50 (35, 60) | 70 (53, 84) | 1.40 | 0.006 |
| Program Promotions | | | | | | |
| Fliers | 37 | 76 | 43 (33, 56) | 69 (55, 80) | 1.60 | 0.005 |
| Email | 34 | 69 | 50 (45, 71) | 66 (54, 84) | 1.32 | 0.05 |
| Consequences upon Declination | | | | | | |
| None | 37 | 76 | 86 (82, 93) | 55 (45, 70) | 0.64 | 0.0001 |
| Some consequence | 12 | 24 | 55 (45, 70) | 86 (82, 93) | 1.56 | 0.0001 |
| Wear mask | 10 | 20 | 56 (46, 70) | 89 (85, 94) | 1.59 | 0.0001 |



Conclusions

- Preliminary results demonstrate large variability among influenza vaccination programs in Louisiana hospitals. No hospital required vaccination as a condition of employment.
- Hospitals that impose consequences for vaccine declination have a higher vaccination rate than hospitals without consequences.
- Our findings suggest that to reach the goal of 90% vaccination rate by 2020, programs with consequences for declination (e.g. wearing a mask) must be enforced.
- These findings have important public health implications.

Examining Louisiana Mammography Facilities for Medicaid Coverage Gaps

Jessica Anderson¹, Courtney Wheeler², Dr. Donna Williams²

Xavier University of Louisiana¹, Louisiana State University School of Public Health, Louisiana Breast and Cervical Health Program²



Introduction


In Louisiana, breast cancer poses the greatest burden of cancer for leading cancer mortality for women (1). Breast cancer also has the highest incidence rate in the state for cancer in women. African-American women have higher incidence and mortality rates than White women, both in the state of Louisiana and the United States (2).

With that being said, it is important for the Louisiana health care system to better understand the state's breast cancer rates in through prevention and diagnostic screening with mammography. The US Preventive Services Task Force (USPSTF) recommends that women over the age of 55 to get a mammogram every 1 year (3). Recently, there have been changes to the Medicaid coverage policy, which extended coverage to more patients in more locations. This coverage includes mammography services. However, there are gaps in coverage across the state of Louisiana, where women cannot acquire adequate care and coverage for mammography and breast screening.

The goal of this research project is to evaluate the patient population of Medicaid-covered, age 55+, who have no access to a breast MRI given at the facility nearest their ZIP. We would like to see where all the mammography sites that accept Medicaid insurance are located, so that we can locate coverage gaps. We will compare the location data to both the uninsured rate and resident rate in the state of Louisiana.





Methods

1. Identified medical facilities providing mammography and breast imaging and created a spreadsheet.
2. Contact each facility to find out information on whether or not the facility accepts Medicaid insurance and, if so, which Medicaid plan was accepted. Information was documented on the spreadsheet.
3. Created a map of the facilities which accept Medicaid insurance in order to see where gaps of coverage and care are located. Geographic information system software (ArcMap) was used to create the coverage map (4).



Results

Maps show that the majority of the facilities are in the Southern part of the state with many being within a 1 or 10 mile radius of the center (Figure 1). There are large clusters of facilities in the Bossier Parish area, the Baton Rouge Parish area, as well as the Lafayette Parish area. The maps also identified gaps in coverage. Bossier Parish locations, which appear to be more rural, have no direct clusters and the facilities are more spread out. While the population of these rural areas tend to be lower than more urban areas, they tend to have higher uninsured rates (Figure 1 and 2). There is no specific trend in the uninsured rate. In all counties, however, there are higher uninsured rates in the more urban, Southern portion of the state in comparison to the Northern part of the state (Figure 2). There are coverage gaps in the lower Southern and Southern portions of the state in Bossier, Morehouse, Rapides, and St. Bernard Parishes. There are also coverage gaps located in the Thibodaux, Iberville, and Terrebonne Parishes areas. Facilities have coverage gaps in Bossier, Rapides, and St. Bernard Parishes (Figure 3). There is a trend of large uninsured rates in the Southern portion of the state, however, there are larger uninsured coverage gaps in the Northern region of the state (Figure 4).



Note in Figures 1, 2, and 3, only mammography facilities that accept Medicaid insurance are shown.

Conclusions

The maps generated revealed some gaps in coverage for the target population of women, age 55+ from an access to a breast MRI while breast cancer care (BPC). There are gaps of coverage in the Thibodaux, Iberville, and Terrebonne Parish areas, as well as the Southern and Southern portions of the state. There is also a trend from Lafayette Parish moving Southwest to Bossier Parish where there is lack of adequate coverage.

When comparing these coverage gaps to the maps displaying French-Speaking Areas, uninsured rates and uninsured rates, we can clearly see where these coverage gaps can be more severe than other areas tend to have higher uninsured rates and large uninsured percentages. For example, in Rapides Parish, there is a large uninsured rate as well as a large uninsured rate and the large uninsured percentage. While the population of women is not very large, the combination of the lack of providers, insurance, and uninsured rates may account for the population of that area. Similarly, Thibodaux and Bossier Parish have coverage gaps. While Parish has a large uninsured rate and does not have any mammography facilities within its borders, the other two lack. Bossier Parish has a lower uninsured rate, but it lacks providers and has a high uninsured percentage. This could vary from its location, they both do not have a proper mammography services.

The state clearly indicates areas where there are gaps of coverage. Some providers do not even have a facility within their borders. In other areas, women may have to travel over 10 miles to reach the nearest mammography facility. It is essential for the state of Louisiana to provide these patients with adequate coverage for mammography services. These coverage gaps may be attributed to the high uninsured and uninsured rates. Closing these gaps would help lower these numbers.

Next Steps

There are a few different options that can be explored when trying to close these coverage gaps. One option is to have a mobile mammography unit travel to a single location in the parish once or twice a month to give mammography. Another option is to create partnerships with different medical facilities across the state to provide mammography for patients with Medicaid insurance. Finally, creating centers can be built in several locations to provide mammography for these patients. After a few years, it would be necessary to analyze the impact of these changes in relation to the uninsured and uninsured rates in the state of Louisiana.

References

1. CDC. • Breast Cancer, Epidemiology, and Control (Breast Cancer Statistics 2020). National Breast Foundation. 2021. <https://www.nbcf.org/breast-cancer-statistics-2020/>. Accessed July 15, 2021.
2. Williams, M., Williams, W., & Bouillon-Bu, A. Louisiana Breast and Cervical Health Program. 2021. <https://www.lsuhs.edu/health-services/health-services/breast-and-cervical-program>. 2021.
3. United States Preventive Services Task Force (USPSTF). (2016). Final Recommendation Statement: Breast Cancer: Screening. Retrieved from <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/breast-cancer-screening>
4. Thomson Reuters (2020). ArcMap Software. Retrieved from <https://www.esri.com/en-us/software/arcmap/arcmap-software/arcmap-software-map-logs>
5. 2018 Census. Available from <https://www.census.gov/data/tables/2018/census-tracts/2018-census-tracts.html>. Environmental Systems Research Institute.

Important Deadlines #4 and 5:

- Record your presentation using Zoom.
- Present your poster for about 5 -7 minutes.
- Send the link to SoMHonorsProgram@lsuhsc.edu by **Monday, October 18th**.
- The judges will receive the abstract book and your pre-recorded presentation on October 22nd, and you will have to answer questions using Zoom.
- Check your email and anticipate about three judges to email you to arrange a 10-15 minute **Zoom meeting between Friday, October 22nd-Tuesday, Oct 26th**. You will have to answer questions regarding your poster.

The presence of soil microbes reduces the allelopathic effects of thymol on tomatoes

Jake Groen, Lexi Smith, Trust Amitaye
Botany Lab BIOL 302L, Science Department



Introduction

Tomatoes (*Solanum lycopersicum*) are often grown close to other plants that may exert a negative effect on the tomatoes through allelopathy, the production of biochemicals that influence other plants success. A garden herb with allelopathic potential is thyme (*Thymus vulgaris*). The objective of this experiment was to determine if the presence of soil microbes inhibits the allelopathic action of thyme's allelochemical, pure white thymol on tomatoes. This information could benefit gardeners, commercial growers, or other researchers.

Grass treated with thyme in the presence of soil microbes had higher survival rates than grass treated with thyme in sterile soil. The microbes helped to overcome thyme's negative effects¹. Bacteria in soil can serve as an important modulator in plant-plant allelopathic interactions².

We hypothesize that the presence of soil microbes will have a positive impact on the germination and growth of *Solanum lycopersicum* when treated with any concentration of thymol as compared to those in sterile soil.

Materials and Methods

Experimental Design

- 8 treatments total: sterilized and non-sterilized soil each treated with four levels of pure white thymol: 0, 12.5 (low), 25 (medium), or 50 uL/100g (high) (3 pots/group)
- Five seeds of tomatoes were placed in each 11.5-centimeter pot
- Plants were grown in greenhouse and separated by cardboard to prevent cross-contamination
- Watered as needed

Soil Preparation

- PRO-MIX soil was steam sterilized for 45 minutes at 125 °C
- 1/4 tsp. osmocote was added to each pot
- Diffusion discs with low, medium, and high concentrations of pure white thymol were applied to each soil treatment, mixed and put into a sealed container to homogenize the thymol and soil

Materials and Methods

Data Collection

- We measured germination percentage and the dry weight at end of the experiment
- Plant material was dried at 60°C for 48 hours

Results – Percent Germination



Figure 1. Experimental set up. Plants grown in sterile or nonsterile soil with no, low, medium, or high levels of thymol.

- Percent germination was highest in sterile soils with no thymol and nonsterile soils at low thymol concentration
- When thymol was present, percent germination was always higher in nonsterile than in sterile soil treatments




Figure 2. Seedling germination percent of tomato seeds treated with zero, low, medium, and high concentrations of pure white thymol in both sterile and non-sterile soil. Mean, ± SE, n=8. All sterile 25uL/100g pots had the same germination %, so the standard error was 0.

Results – Biomass

- When thymol was present, biomass was greater in sterile than in non-sterile soil treatments
- The mean value difference in 12.5 uL/100g and 25 uL/100g were **not significant/small**.
- Control groups showed that nonsterile soil had a significantly greater biomass at 0.0924g while sterile soil had a 0.0411g biomass.

Average Above Ground Biomass




Figure 3. Above ground biomass of tomato seeds treated with zero, low, medium, and high concentrations of pure white thymol in both sterile and non-sterile soil. Means ± SE, n = 8.

Discussion

The germination results supported our hypothesis, but the biomass findings were mixed. As hypothesized, all seeds grown in nonsterile soil and treated with thymol showed a higher germination percentage than seeds grown in sterile soil at the same thymol levels. In contrast, the biomass of plants grown in sterile soil and treated with thymol was higher than that of plants grown in nonsterile soil at the corresponding levels of thymol, though the differences were small at low and medium thymol levels. When no thymol was applied, germination and biomass showed opposite patterns: germination was higher in sterile soils, but biomass was higher in nonsterile soils.

Discussion (cont.)

Our germination results support other studies that found soil microbes degrade the allelopathic effects of thymol into a nonharmful substance for tomatoes¹. Our biomass results contradict other studies that show how microbes metabolize toxic chemicals, so the plant's growth is not inhibited by them². Overall, the results suggest that microbes aid in germination, but don't provide extra benefit for later life stages at high thymol concentrations.

Limitations

Early harvesting is a limitation of our experiment. The biomass results may have been skewed by the plants that germinated first. The tomatoes may not have reached their full potential before they were harvested. **Another limitation may be old tomato seeds.**

Conclusions

These results support a diverse and healthy social microbial community. One recommendation would be to avoid intercropping thyme with tomatoes. Further research could examine food crops from the nightshade family to determine how widespread the effects of thyme allelopathy are. **This experiment could be repeated and carried out its full course.**

Literature Cited

- Ehlers, B. K. (2011). Soil Microorganisms Alleviate the Allelochemical Effects of a Thymus Monoterpene on the Performance of an Associated Grass Species. *PLoS One*, 6(11).
- Milner, B. (2015). Unravelling the beneficial role of microbial contributors in reducing the allelopathic effects of weeds. *Applied Microbiology and Biotechnology*, 97(12), 5050-5055.


Acknowledgments

Thank you to Professor Flint for providing additional knowledge and advice throughout the experiment. Thanks to Jacky Alesen and Bailey Dell for assistance with watering our plants in the greenhouse. Lastly, thanks to Dr. Greenfield for supplying us with thymol and diffusion discs.




Important deadline Number 6:

- The Poster Symposium Awards ceremony will be held in person on **October 29th, 2:00-2:30 PM**
- Students, mentors, and lab personnel may attend in person.
- Families are welcome to view the live presentation via Zoom.
- Keep in mind that these plans may change due to changing COVID-19 guidelines.



Next: Practice your presentation


- Practice with your mentor and your lab members!
 - Anticipate questions and look up the answers ahead of time
 - Practice, practice, practice so you sound polished.
 - Practice in front of your friends or in front of a mirror.
- 

What happens at a virtual poster session?

- Please dress appropriately (business attire or lab coat) during the recorded presentation and meeting with the judges
- The abstracts and recorded posters will be sent to the judges on October 22nd.
- The posters will be judged on the actual poster display and your presentation (enthusiasm, understanding of the topic, etc)



What happens at a virtual poster session?

- Sometimes people ask specific questions, or sometimes they ask “tell me about your project”
 - Think of possible questions you may be asked. If you do not know an answer, it is OK to say “I don’t know”
 - The abstracts and posters will be available to the public on October 22nd .
- 




Zoom Presentation example:



Molecular Cytogenetic Characterization of RH4 and RH30 Alveolar Rhabdomyosarcoma (ARMS) Cell Lines

Jorge Peñas¹, Katrina Gleditsch^{1,2}, Danielle Mercer¹,
Ayesha Umrigar¹, Yuwen Li³, Tian-Jian Chen³, Andrew
Hollenbach¹, Fern Tsien¹



¹ Louisiana State University Health Sciences Center, Department of Genetics, ²
Children's Hospital of New Orleans, ³Tulane School of Medicine Hayward Genetics
Center

Molecular Cytogenetic Characterization of RH4 and RH30 Alveolar Rhabdomyosarcoma (ARMS) Cell Lines



Jorge Peñas¹, Katrina Gleditsch^{1,2}, Danielle Mercer¹, Ayesha Umrigar¹, Yuwen Li³, Tian-Jian Chen³, Andrew Hollenbach¹, Fern Tsien¹

¹Louisiana State University Health Sciences Center, Department of Genetics, ²Children's Hospital of New Orleans, ³Tulane School of Medicine Hayward Genetics Center


Background

Rhabdomyosarcoma is the most common soft tissue sarcoma in children. More than half of childhood rhabdomyosarcomas are diagnosed in those under the age of 10. The two most common subtypes of rhabdomyosarcoma are alveolar (ARMS) and embryonal (ERMS).

- ERMS makes up 60% of rhabdomyosarcoma cases
- ARMS makes up 30% of cases and typically has a more aggressive clinical course and portends a poorer clinical prognosis.
- Details on ARMS:
 - Historiologically distinct
 - Genetically characterized by balanced translocations at t(2;13) or t(1;13).
 - Translocations lead to the production of fusion proteins PAX3-FOXO1 t(2;13) and PAX7-FOXO1 t(1;13)
 - PAX3-FOXO1 variant of ARMS is known to be more aggressive.
 - Current diagnostic break apart probe cannot distinguish between translocations



Our laboratory, in collaboration with Cytocell Ltd., designed fluorescence *in situ* hybridization (FISH) probes specific for the PAX3-FOXO1 and PAX7-FOXO1 translocations, which allows for a more accurate diagnosis by differentiating between t(1;13) and t(2;13).



Two of the most widely used commercially available cell lines used in ARMS research are RH4 and RH30. These cell lines were derived from pediatric patients clinically diagnosed with ARMS.

| Characteristic | RH4 | RH30 |
|----------------|-----------------|--------------|
| Age | 7 Years Old | 24 Years Old |
| Sex | Female | Male |
| Origin | Lung metastasis | Bone Marrow |
| Gene Fusion | t(2;13) | t(2;13) |

Previous studies have cytogenetically characterized the RH30 cell line to gain a better understanding of phenotypic clinical correlations due to these translocations; however, to date the RH4 cell line has not been fully cytogenetically characterized.

The specific aims of this project are to 1) cytogenetically characterize the RH4 and RH30 cell lines since these cell lines are used in clinical research where cytogenetic variants can affect clinical outcomes and to 2) validate our newly designed ARMS probes.

These experiments are also being performed on patient derived ARMS tumor samples thus further validating the newly designed probes to determine their clinical diagnostic utility. (See Gleditsch, et al. poster #2818)

Methods


Table 2: Description and brief protocol of methods used to validate the newly designed probes

| Method | Protocol |
|--|---|
| aCGH+SNP | Array CGH was performed on SurePrint G3 Human CGH+SNP 4x1.80k array slides, scanned by Agilent DNA Microarray Scanner with SureScan High-Resolution, and analyzed by Agilent Feature Extraction and Cytogenomics. |
| G-Banding (Giemsa Banding) | Chromosomes were harvested from RH4, RH30 and normal cells using colcemid, 0.075 M KCl, and Carnoy's fixative (3:1 Methanol: Glacial Acetic Acid). Slides were G-banded using trypsin and Giemsa stain. Slides were analyzed at 100X using a bright field microscope. |
| SKY (Spectral Karyotyping) | Chromosomes were prepared as above. SKY was performed using 24 chromosome-specific probes (Molecular Cytogenetics Core Facility, MD Anderson Cancer Center) and analyzed at 100X with a fluorescent microscope. |
| FISH (Fluorescence <i>in situ</i> Hybridization) | Chromosomes were prepared as above and hybridized with newly designed probes specific for PAX3, FOXO1, and PAX7. These probes were labeled with Spectrum Orange, FITC, & Spectrum Aqua, respectively, followed with a DAPI counterstain, and analyzed at 100X using a fluorescent microscope. |

Results

FISH

Figure 1: ARMS probes used on RH4 and RH30 cell lines.




Karyotype G-banding

Figure 2: Karyotype (Left: RH4, Right: RH30)



Ploidy of RH4 and RH30 G-Banded Cells



SKY (RH4)




Figure 3: Spectral karyotyping of one representative cell from RH4 cell line. A) Metaphase spread B) Organized karyotype demonstrates aneuploidy and multiple chromosomal rearrangements. Up to 5 rearrangements are shown in a single chromosome. Each chromosome is displayed in triplicate: G-Banded, fluorescent, and a computer assigned color, respectively.

aCGH

Figure 4: arrayCGH and expanded regions of chromosome 17.




Table 2: Copy number variations (CNV) between RH4 and RH30 cell lines determined by array Comparative Genomic Hybridization (arrayCGH).

| Gene | Location | Function | RH4 | RH30 |
|-------|----------|---|-----|------|
| WRAS | 1p13.2 | Provides instructions for making a protein that is involved primarily in regulating cell division. | + | 0 |
| MYCN | 2p24.1 | Proto-oncogene member of the MYC family of transcription factors. | 0 | + |
| MET | 7q31.2 | Mesenchymal Epithelial Transition MET is a prototypical receptor tyrosine kinase. | 0 | + |
| CDK4 | 12q13.3 | Involved in cell cycle progression. | 0 | + |
| TP53 | 17p13.1 | Codes for a protein that regulates the cell cycle and functions as a tumor suppressor. | + | 0 |
| MDM2 | 12q14.3 | Encodes a nuclear-localized E3 ubiquitin ligase. Target tumor suppressor proteins, such as p53. | + | 0 |
| PAX3 | 2q35 | These genes play critical roles during fetal development. Necessary for myogenesis. | + | 0 |
| PAX7 | 1p36.13 | Transcription factor playing a role in myogenesis through regulation of muscle precursor cells proliferation. | + | 0 |
| FOXO1 | 13q14.1 | forkhead family of transcription factors. Play a role in myogenic growth and differentiation. | 0 | + |

Discussion

This study provides the cytogenetic and molecular genetic characterization of commonly used ARMS cell lines, RH4 and RH30. The cell lines were analyzed using G banding, fluorescence *in situ* hybridization (FISH), array comparative genomic hybridization (aCGH) and spectral karyotyping (SKY).

FISH results using our newly developed probes specific for PAX3, PAX7, and FOXO1 showed that RH30 had more PAX3-FOXO1 fusions when compared to RH4. However, RH4 had a significant amplification of PAX3, PAX7, and FOXO1.

Consistent with our FISH results, G-Banding demonstrated a significant difference in ploidy between RH4 and RH30. Aneuploidy for RH4 ranged from 59-163 chromosomes, whereas RH30 had 51-90 chromosomes.

Initial SKY analysis of RH4 shows large scale and highly complex chromosomal rearrangements and aneuploidy throughout the genome.

ArrayCGH showed similarities and differences in chromosomal amplifications and deletions between the two cell lines. Genes within similar regions were selected from previously published reports with known association to rhabdomyosarcoma. The amplification of PAX3 in RH4 seen in our aCGH is consistent with our FISH results, which serves as an internal positive control.

While both cell lines originate from patients diagnosed with ARMS, they exhibit significant cytogenetic variations, highlighting the necessity of genetic characterization to direct potential targeted treatment options.

Our future aims for this study is to further characterize the genetic variation among ARMS cell lines and patient tumors with bioinformatics and pathways analysis software. Furthermore, RNA sequencing and epigenetic analysis will be performed on ARMS tumor samples.


Currently the second arm of our study is looking at the heterogeneity of 2;13 translocations in ARMS patients at hospitals in New Orleans further validating our newly designed FISH probes. (See Gleditsch, et al. poster #2818)



Background

Rhabdomyosarcoma is the most common soft tissue sarcoma in children. More than half of childhood rhabdomyosarcomas are diagnosed in those under the age of 10. The two most common subtypes of rhabdomyosarcoma are alveolar (ARMS) and embryonal (ERMS).


- ERMS makes up 60% of rhabdomyosarcoma cases
- ARMS makes up 30% of cases and typically has a more aggressive clinical course and portends a poorer prognosis.
- Details on ARMS:
 - Histologically distinct
 - Genetically characterized by balanced translocations at t(2;13) or t(1;13).
 - Translocations lead to the production of fusion proteins PAX3-FOXO1 t(2;13) and PAX7-FOXO1 t(1;13)
 - PAX3-FOXO1 variant of ARMS is known to be more aggressive.
 - Current diagnostic break apart probe cannot distinguish between translocations



t(2;13)(q35;q14)

PAX3 – FOXO1 Fusion


More Aggressive,
Worse Prognosis



t(1;13)(p36;q14)

PAX7 – FOXO1 Fusion

Our laboratory, in collaboration with Cytocell Ltd., designed fluorescence *in situ* hybridization (FISH) probes specific for the PAX3-FOXO1 and PAX7-FOXO1 translocations, which allows for a more accurate diagnosis by differentiating between t(1;13) and t(2;13).



Two of the most widely used commercially available cell lines used in ARMS research are RH4 and RH30. These cell lines were derived from pediatric patients clinically diagnosed with ARMS.

Table 1: Characteristics of RH4 and RH30 cell lines

| | RH4 | RH30 |
|--------------------|-----------------|--------------|
| Age | 7 Years Old | 16 Years Old |
| Sex | Female | Male |
| Origin | Lung metastasis | Bone Marrow |
| Gene Fusion | t(2:13) | t(2:13) |

Previous studies have cytogenetically characterized the RH30 cell line to gain a better understanding of phenotypic clinical correlations due to these translocations; however, to date the RH4 cell line has not been fully cytogenetically characterized.

The specific aims of this project are to 1) cytogenetically characterize the RH4 and RH30 cell lines since these cell lines are used in clinical research where cytogenetic variants can affect clinical outcomes and to 2) validate our newly designed ARMS probes.

These experiments are also being preformed on patient derived ARMS tumor samples thus further validating the newly designed probes to determine their clinical diagnostic utility. (**See Gleditsch, *et al.* poster #2818**)

Methods

aCGH+SNP (array Comparative Genomic Hybridization + Single Nucleotide Polymorphism)

Array CGH was performed on SurePrint G3 Human CGH+SNP 4x180k array slides, scanned by Agilent DNA Microarray Scanner with SureScan High-Resolution, and analyzed by Agilent FeatureExtraction and Cytogenomics.

G-Banding (Giemsa Banding)

Chromosomes were harvested from RH30, RH4 and normal cells using colcemid, 0.075 M KCl, and Carnoy's fixative (3:1 Methanol: Glacial Acetic Acid). Slides were G-banded using trypsin and Giemsa stain. Slides were analyzed at 100X using a bright field microscope.

SKY (Spectral Karyotyping)

Chromosomes were prepared as above. SKY was performed using 24 chromosome-specific probes (Molecular Cytogenetics Core Facility, MD Anderson Cancer Center) and analyzed at 100X with a fluorescent microscope.


FISH (Fluorescence *in situ* Hybridization)

Chromosomes were prepared as above and hybridized with newly designed probes specific for PAX3, FOXO1, and PAX7. These probes were labeled with Spectrum Orange, FITC, & Spectrum Aqua, respectively, followed with a DAPI counterstain, and analyzed at 100X using a fluorescent microscope.


Results

FISH


Figure 1: ARMS probes used on RH4 and RH30 cell lines.





A) RH4 cell hybridized with PAX3 & FOXO1 probes. Fusions pointed out by arrow. (3 Shown)



B) RH30 cell hybridized with PAX3 & FOXO1 probes. Fusions pointed out by arrow. (4 Shown)






C) RH4 cell hybridized with PAX7 & FOXO1 probes. Images demonstrate amplification of PAX7 and FOXO1 in both cell lines.

D) RH30 PAX7 & FOXO1 probes. Images demonstrate amplification of PAX7 and FOXO1 in both cell lines.

PAX3, PAX7 & FOXO1 signal comparison between RH4 and RH30




Karyotype G-banding

Figure 2: Karyotype (Left: RH4, Right: RH30)



Ploidy of RH4 and RH30 G-Banded Cells



SKY (RH4)






Figure 3: Spectral karyotyping of one representative cell from RH4 cell line. **A)** Metaphase spread **B)** Organized karyotype demonstrates aneuploidy and multiple chromosomal rearrangements. Up to 5 rearrangements are shown in a single chromosome. Each chromosome is displayed in triplicate: G-Banded, fluorescent, and a computer assigned color, respectively.



aCGH

Figure 4: arrayCGH and expanded regions of chromosome 17.



ArrayCGH of **A)** RH4 and **B)** RH30 cells. Expanded regions of Chromosome 17. Purple box highlights 17p13.1 (TP53) in **C)** RH4 and **D)** RH30

Table 2: Copy number variations (CNV) between RH4 and RH30 cell lines determined by array Comparative Genomic Hybridization (arrayCGH).

| Gene | Location | Function | RH4 | RH 30 |
|--------------|-----------------|---|------------|--------------|
| NRAS | 1p13.2 | Provides instructions for making a protein that is involved primarily in regulating cell division. | + | 0 |
| MYCN | 2p24.1 | Proto-oncogene member of the MYC family of transcription factors. | 0 | + |
| MET | 7q31.2 | Mesenchymal Epithelial Transition MET is a prototypical receptor tyrosine kinase. | 0 | + |
| CDK4 | 12q13.3 | Involved in cell cycle progression. | 0 | + |
| TP53 | 17p13.1 | Codes for a protein that regulates the cell cycle and functions as a tumor suppressor. | - | - |
| MDM2 | 12q14.3 | Encodes a nuclear-localized E3 ubiquitin ligase. Target tumor suppressor proteins, such as p53. | + | 0 |
| PAX3 | 2q35 | These genes play critical roles during fetal development. Necessary for myogenesis. | + | 0 |
| PAX7 | 1p36.13 | Transcription factor playing a role in myogenesis through regulation of muscle precursor cells proliferation. | + | - |
| FOXO1 | 13q14.1 | Forkhead family of transcription factors. Play a role in myogenic growth and differentiation. | 0 | + |

Discussion

This study provides the cytogenetic and molecular genetic characterization of commonly used ARMS cell lines, RH4 and RH30. The cell lines were analyzed using G banding, fluorescence *in situ* hybridization (FISH), array comparative genomic hybridization (aCGH) and spectral karyotyping (SKY).

FISH results using our newly developed probes specific for PAX3, PAX7, and FOXO1 showed that RH30 had more PAX3-FOXO1 fusions when compared to RH4. However, RH4 had a significant amplification of PAX3, PAX7, and FOXO1.

Consistent with our FISH results, G-Banding demonstrated a significant difference in ploidy between RH4 and RH30. Aneuploidy for RH4 ranged from 59-163 chromosomes, whereas RH30 had 51-90 chromosomes.

Initial SKY analysis of RH4 shows large scale and highly complex chromosomal rearrangements and aneuploidy throughout the genome.

ArrayCGH showed similarities and differences in chromosomal amplifications and deletions between the two cell lines. Genes within similar regions were selected from previously published reports with known association to rhabdomyosarcoma. The amplification of PAX3 in RH4 seen in our aCGH is consistent with our FISH results, which serves as an internal positive control.

While both cell lines originate from patients diagnosed with ARMS, they exhibit significant cytogenetic variations, highlighting the necessity of genetic characterization to direct potential targeted treatment options.



Our future aims for this study is to further characterize the genetic variation among ARMS cell lines and patient tumors with bioinformatics and pathways analysis software. Furthermore, RNA sequencing and epigenetic analysis will be preformed on ARMS tumor samples.

Currently the second arm of our study is looking at the heterogeneity of 2;13 translocations in ARMS patients at hospitals in New Orleans further validating our newly designed FISH probes. (See Gleditsch, *et al.* poster #2818)

- Practice with your mentor and lab mates.
- After you create your final PowerPoint presentation, record yourself presenting it using Zoom.
- Before recording, do not save in the computer hard drive; save it on the Cloud
- Refer to the 2020 Symposium:
https://www.medschool.lsuhschool.edu/genetics/2020_medical_student_research_virtual_poster_symposium.aspx

Awards :

- Travel awards to present your project at a national conference



Remember :

1. Notification of participation: Friday, September 24th
2. Abstracts due: Monday, October 4th
3. Posters due: Wednesday, October 13th
4. Recording due: Monday, October 18th
5. Zoom appointments with three judges:
Friday, October 22nd-Tuesday, Oct 26th (time: judges will arrange with students)
6. Virtual Poster Symposium Awards :
October 29th, 2:00-2:30 PM

Send the above to: SoMHonorsProgram@lsuhsc.edu

Refer to the 2020 Symposium:

https://www.medschool.lsuhschool.edu/genetics/2020_medical_student_research_virtual_poster_symposium.aspx